

VIRAL POLYMERASE INHIBITORS

RELATED APPLICATIONS

Benefit of US Provisional Application, Serial No. 60/441,674, filed on January 22,
5 2003, is hereby claimed and said application is herein incorporated by reference in
its entirety.

TECHNICAL FIELD OF THE INVENTION

The invention relates to inhibitors of RNA dependent RNA polymerases, particularly
10 those viral polymerases within the Flaviviridae family, more particularly to HCV
polymerase.

BACKGROUND OF THE INVENTION

About 30,000 new cases of hepatitis C virus (HCV) infection are estimated to occur
15 in the United States each year (Kolykhalov, A.A.; Mihalik, K.; Feinstone, S.M.; Rice,
C.M.; 2000; *J. Virol.* **74**: 2046-2051). HCV is not easily cleared by the hosts'
immunological defences; as many as 85% of the people infected with HCV become
chronically infected. Many of these persistent infections result in chronic liver
disease, including cirrhosis and hepatocellular carcinoma (Hoofnagle, J.H.; 1997;
20 *Hepatology* **26**: 15S-20S*). There are an estimated 170 million HCV carriers world-
wide, and HCV-associated end-stage liver disease is now the leading cause of liver
transplantation. In the United States alone, hepatitis C is responsible for 8,000 to
10,000 deaths annually. Without effective intervention, the number is expected to
triple in the next 10 to 20 years. There is no vaccine to prevent HCV infection.
25 Prolonged treatment of chronically infected patients with interferon or interferon and
ribavirin is the only currently approved therapy, but it achieves a sustained response
in fewer than 50% of cases (Lindsay, K.L.; 1997; *Hepatology* **26**: 71S-77S*, and
Reichard, O.; Schvarcz, R.; Weiland, O.; 1997 *Hepatology* **26**: 108S-111S*).

30 HCV belongs to the family *Flaviviridae*, genus *hepacivirus*, which comprises three
genera of small enveloped positive-strand RNA viruses (Rice, C.M.; 1996;
"Flaviviridae: the viruses and their replication"; pp. 931-960 in *Fields Virology*; Fields,
B.N.; Knipe, D.M.; Howley, P.M. (eds.); Lippincott-Raven Publishers, Philadelphia
Pa. *). The 9.6 kb genome of HCV consists of a long open reading frame (ORF)
35 flanked by 5' and 3' non-translated regions (NTR's). The HCV 5' NTR is 341

nucleotides in length and functions as an internal ribosome entry site for cap-independent translation initiation (Lemon, S.H.; Honda, M.; 1997; *Semin. Virol.* **8**: 274-288). The HCV polyprotein is cleaved co- and post-translationally into at least 10 individual polypeptides (Reed, K.E.; Rice, C.M.; 1999; *Curr. Top. Microbiol. Immunol.* **242**: 55-84*). The structural proteins result from signal peptidases in the N-terminal portion of the polyprotein. Two viral proteases mediate downstream cleavages to produce non-structural (NS) proteins that function as components of the HCV RNA replicase. The NS2-3 protease spans the C-terminal half of the NS2 and the N-terminal one-third of NS3 and catalyses *cis* cleavage of the NS2/3 site. The same portion of NS3 also encodes the catalytic domain of the NS3-4A serine protease that cleaves at four downstream sites. The C-terminal two-thirds of NS3 is highly conserved amongst HCV isolates, with RNA-binding, RNA-stimulated NTPase, and RNA unwinding activities. Although NS4B and the NS5A phosphoprotein are also likely components of the replicase, their specific roles are unknown. The C-terminal polyprotein cleavage product, NS5B, is the elongation subunit of the HCV replicase possessing RNA-dependent RNA polymerase (RdRp) activity (Behrens, S.E.; Tomei, L.; DeFrancesco, R.; 1996; *EMBO J.* **15**: 12-22*; and Lohmann, V.; Körner, F.; Herian, U.; Bartenschlager, R.; 1997; *J. Virol.* **71**: 8416-8428*). It has been recently demonstrated that mutations destroying NS5B activity abolish infectivity of RNA in a chimp model (Kolykhalov, A.A.; Mihalik, K.; Feinstone, S.M.; Rice, C.M.; 2000; *J. Virol.* **74**: 2046-2051*).

The development of new and specific anti-HCV treatments is a high priority, and virus-specific functions essential for replication are the most attractive targets for drug development. The absence of RNA dependent RNA polymerases in mammals, and the fact that this enzyme appears to be essential to viral replication, would suggest that the NS5B polymerase is an ideal target for anti-HCV therapeutics. WO 00/06529, WO 00/13708, WO 00/10573, WO 00/18231, WO 01/47883, WO 01/85172, WO 02/04425, WO 03/010140 and WO 03/010141 report inhibitors of NS5B proposed for treatment of HCV.


SUMMARY OF THE INVENTION

It is therefore an object of the invention to provide a novel series of compounds having inhibitory activity against HCV polymerase.

Further objects of this invention arise for the one skilled in the art from the following description and the examples.

In a first aspect of the invention, there is provided an isomer, enantiomer,

- 5 diastereoisomer or tautomer of a compound, represented by formula I:



wherein

10

----- represents either a single or a double bond;

B is -N- and **A** is =CR¹- or =N-; or

- 15 **B** is =C- and **A** is O, S or NR¹;

R¹ is selected from the group consisting of: H, (C₁₋₆)alkyl optionally substituted with:

- 20 halogen, OR¹¹, SR¹¹ or N(R¹²)₂, wherein **R**¹¹ and each **R**¹² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-Het, said aryl or Het optionally substituted with **R**¹⁶⁰; or both **R**¹² are covalently bonded together and to the nitrogen to which they are both attached to form a 5, 6 or 7-membered saturated heterocycle;

- 25 the group -C(=Y¹)-Z is covalently linked to either **M**² or **M**³,

M¹ is CR^{4a},

M² or **M**³, when not linked to -C(=Y¹)-Z, is CR⁵,

M⁴ is CR^{4b},

30

and in addition one or two of the groups selected from **M**¹, **M**², **M**³ and **M**⁴ may also

be N, with the proviso that the group M^2 or M^3 to which $-C(=Y^1)-Z$ is linked is a C-atom,

Y^1 is O or S;

5

Z is defined as $NR^{N2}-SO_2-R^C$ or $NR^{N3}-SO_2-N(R^{N2})R^{N1}$, wherein R^C , R^{N1} or any heterocycle formed by R^{N1} and R^{N2} is optionally substituted with R^{60} ;

10 R^2 is selected from: halogen or R^{21} , wherein R^{21} is aryl or Het, said R^{21} is optionally substituted with R^{150} ;

R^3 is selected from (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl, (C₅₋₇)cycloalkenyl, (C₁₋₃)alkyl-(C₅₋₇)cycloalkenyl, (C₆₋₁₀)bicycloalkyl, (C₁₋₃)alkyl-(C₆₋₁₀)bicycloalkyl, (C₆₋₁₀)bicycloalkenyl, (C₁₋₃)alkyl-(C₆₋₁₀)bicycloalkenyl, **HCy** or (C₁₋₃)alkyl-**HCy**,

15

wherein **HCy** is a saturated or unsaturated 4 to 7-membered heterocyclic group with 1 to 3 heteroatoms selected from O, S and N;

said alkyl, cycloalkyl, cycloalkenyl, bicycloalkyl, bicycloalkenyl, **HCy** and alkyl-**HCy** being optionally substituted with from 1 to 4 substituents selected from: a) halogen;

20

b) (C₁₋₆)alkyl optionally substituted with:

- 1 to 3 substituents selected from halogen;

- OR^{31} or SR^{31} wherein R^{31} is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl; or

25

- $N(R^{32})_2$ wherein each R^{32} is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl; or both R^{32} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

30

c) OR^{33} or SR^{33} wherein R^{33} is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl;

d) $N(R^{35})_2$ wherein each R^{35} is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl; or both R^{35} are covalently bonded together and to the nitrogen to which they are attached to

form a 5, 6 or 7-membered saturated heterocycle;

R^{4a} , R^{4b} , R^5 each are independently H or defined as R^{150} ;

- 5 R^{60} is defined as 1 to 4 substituents independently selected from:
- 1 to 3 substituents selected from halogen;
 - one of each substituent selected from: OPO_3H , NO_2 , cyano, azido, $C(=NH)NH_2$, $C(=NH)NH(C_{1-6})alkyl$ or $C(=NH)NHCO(C_{1-6})alkyl$, SO_3H ; and
 - 1 to 3 substituents selected from:
- 10 a) $(C_{1-6})alkyl$, $(C_{3-7})cycloalkyl$, $(C_{3-7})spirocycloalkyl$ optionally containing 1 or 2 heteroatoms selected from N, O and S; $(C_{2-6})alkenyl$, $(C_{2-8})alkynyl$, $(C_{1-6})alkyl-(C_{3-7})cycloalkyl$, all of which optionally being substituted with R^{150} ;
- b) OR^O ;
- 15 c) $OC(O)R^O$;
- d) SR^O , SO_2R^C , $SO_2N(R^{N2})R^{N1}$, $SO_2N(R^{N2})C(O)R^C$, $CONR^{N3}SO_2N(R^{N2})R^{N1}$, or $CONR^{N2}SO_2R^C$;
- e) $N(R^{N2})R^{N1}$, $N(R^{N2})COOR^C$, $N(R^{N2})SO_2R^C$ or $N(R^{N1})OR^O$;
- f) $N(R^{N2})COR^C$;
- 20 g) $N(R^{N3})CON(R^{N2})R^{N1}$;
- h) $N(R^{N3})COCOR^C$, $N(R^{N3})COCOOR^O$, $N(R^{N3})COCON(R^{N2})OR^O$, or $N(R^{N3})COCON(R^{N2})R^{N1}$;
- i) COR^O ;
- j) $COOR^O$;
- 25 k) $CON(R^{N2})R^{N1}$;
- l) aryl, Het, $(C_{1-4})alkyl-aryl$ or $(C_{1-4})alkyl-Het$, all of which optionally being substituted with R^{150} ;

wherein said R^{N1} , R^C and/or R^O are optionally substituted with R^{150} as defined,

30

R^{150} is defined as 1 to 4 substituents independently selected from:

- 1 to 3 substituents selected from halogen;
- one of each substituent selected from: OPO_3H , NO_2 , cyano, azido, SO_3H , $C(=NH)NH_2$, $C(=NH)NH(C_{1-6})alkyl$ or $C(=NH)NHCO(C_{1-6})alkyl$; and

- 1 to 3 substituents selected from:

- a) (C₁₋₆) alkyl, (C₃₋₇)cycloalkyl, (C₃₋₇)spirocycloalkyl optionally containing 1 or 2 heteroatoms selected from N, O and S; (C₂₋₆)alkenyl, (C₂₋₈)alkynyl, (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl, all of which optionally substituted with R¹⁶⁰;
- 5 b) OR⁰;
- c) OC(O)R⁰;
- d) SR⁰, SO₂R^C, SO₂N(R^{N2})R^{N1} or SO₂N(R^{N2})C(O)R^C;
- e) N(R^{N2})R^{N1}, N(R^{N2})COOR^C, N(R^{N2})SO₂R^C or N(R^{N1})OR⁰;
- f) N(R^{N2})COR^C;
- 10 g) N(R^{N3})CON(R^{N2})R^{N1};
- h) N(R^{N3})COCOR^C, N(R^{N3})COCOOR⁰, N(R^{N3})COCON(R^{N2})OH, N(R^{N3})COCON(R^{N2})O(C₁₋₄)alkyl or N(R^{N3})COCON(R^{N2})R^{N1};
- i) COR⁰;
- j) COOR⁰;
- 15 k) tetrazole, triazole, CONR^{N2}SO₂R^C, CONR^{N3}-SO₂N(R^{N2})R^{N1} or CON(R^{N2})R^{N1};

wherein said R^{N1}, R^C and/or R⁰ are optionally substituted with R¹⁶⁰ as defined;

20 R¹⁶⁰ is defined as 1, 2 or 3 substituents independently selected from:

- 1, 2 or 3 fluorine substituents; and
- one of each substituent selected from tetrazole, triazole, chlorine, bromine, iodine, CN, nitro, (C₁₋₄)alkyl, OCF₃, SCF₃, CF₃, COOR¹⁶¹, SO₃H, SR¹⁶¹, SO₂R¹⁶³, OR¹⁶¹, N(R¹⁶²)₂, SO₂N(R¹⁶²)₂, SO₂NR¹⁶²COR¹⁶², NR¹⁶²SO₂R¹⁶³, -NR¹⁶¹-CO-COOR¹⁶¹, -NR¹⁶¹-CO-CO(NR¹⁶²)₂, -CONR¹⁶¹SO₂R^C, CONR¹⁶¹-SO₂N(R¹⁶²)₂ or -SO₂-NR¹⁶¹-COR^C, NR¹⁶²COR¹⁶² or CON(R¹⁶²)₂, wherein R¹⁶¹, R¹⁶³ and each R¹⁶² is independently (C₁₋₄)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₇)cycloalkyl; and R¹⁶¹ and each R¹⁶² may each independently also be H; or both R¹⁶² are covalently bonded together and to the nitrogen to which they
- 25
- 30 are attached to form a 5, 6 or 7-membered saturated heterocycle;

R⁰, R^C are independently defined as (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₄)alkyl-(C₃₋₇)cycloalkyl, (C₂₋₆)alkenyl, aryl, Het, (C₁₋₄)alkyl-aryl, or (C₁₋₄)alkyl-Het; or R⁰ is also optionally defined as H.

R^{N1} is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₄)alkyl-(C₃₋₇)cycloalkyl, (C₂₋₆)alkenyl, aryl, Het, (C₁₋₄)alkyl-aryl, (C₁₋₄)alkyl-Het; and

- 5 R^{N2} , R^{N3} , R^{N4} are independently H, CH₃, (C₂₋₆)alkyl, (C₃₋₆)cycloalkyl, (C₁₋₄)alkyl-(C₃₋₆)cycloalkyl; all of which being optionally substituted with halogen, carboxy or (C₁₋₆)alkoxycarbonyl; and/or wherein said alkyl, cycloalkyl or alkylcycloalkyl is optionally substituted with hydroxy, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, amino, -NH(C₁₋₄)alkyl and/or -N((C₁₋₄)alkyl)₂; or

10

in the case

- a) of a group N(R^{N2})(R^{N1}) the substituents R^{N2} and R^{N1} ; or
b) of a group NR^{N3}-N(R^{N2})(R^{N1}) the substituents R^{N3} and R^{N1} , or R^{N2} and R^{N1} ; may be covalently bonded together to form a 4-, 5-, 6- or 7-membered
15 saturated or unsaturated N-containing heterocycle or a 8-, 9-, 10- or 11-membered N-containing heterobicycle, each optionally having additionally from 1 to 3 heteroatoms selected from O, N, and S, wherein said heterocycle or heterobicycle is optionally substituted as defined;

- 20 wherein Het is defined as a 4-, 5-, 6- or 7-membered heterocycle having 1 to 4 heteroatoms selected from O, N and S, or a 8-, 9-, 10- or 11-membered heterobicycle having 1 to 5 heteroatoms selected from O, N and S;

or a salt thereof.

25

Included within the scope of this invention are compounds of the formula (I) as described hereinbefore, to which a "detectable label", "affinity tag" or "photoreactive group" is linked.

- 30 The compounds according to this invention generally show an inhibitory activity against HCV polymerase. In particular compounds according to this invention inhibit RNA synthesis by the RNA dependent RNA polymerase of HCV, especially of the enzyme NS5B encoded by HCV. Furthermore, compounds being active in cell culture are provided. A further advantage of compounds provided by this invention is
35 their low to very low or even non-significant activity against other polymerases.

In a second aspect of the invention, there is provided a use of a compound of formula I according to this invention, or a pharmaceutically acceptable salt thereof, as an HCV polymerase inhibitor.

5

In a third aspect of the invention, there is provided a use of a compound of the formula I according to this invention, or a pharmaceutically acceptable salt thereof, as an inhibitor of RNA dependent RNA polymerase activity of the enzyme NS5B, encoded by HCV.

10

In a fourth aspect of the invention, there is provided a use of a compound of the formula I according to this invention, or a pharmaceutically acceptable salt thereof, as an inhibitor of HCV replication.

15 In a fifth aspect of the invention, there is provided a method of treating or preventing HCV infection in a mammal, comprising administering to the mammal an effective amount of a compound of formula I according to this invention, or a pharmaceutically acceptable salt thereof.

20 In a sixth aspect of the invention, there is provided a method of treating or preventing HCV infection in a mammal, comprising administering to the mammal an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof in combination with another antiviral agent.

25 In a seventh aspect of the invention, there is provided a pharmaceutical composition for the treatment or prevention of HCV infection, comprising an effective amount of a compound of formula I according to this invention, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

30 According to a further specific embodiment, the pharmaceutical composition of this invention comprises a therapeutically effective amount of one or more antiviral agents. Examples of antiviral agents include, ribavirin and amantadine.

35 According to a further specific embodiment, the pharmaceutical composition of this invention comprises an other anti-HCV agent as an antiviral agent.

According to a more specific embodiment, the pharmaceutical composition of this invention comprises an additional immunomodulatory agent as an other anti-HCV agent. Examples of additional immunomodulatory agents include but are not limited to, α -, β -, δ -, γ -, tau- and ω -interferons.

According to another more specific embodiment, the pharmaceutical composition of this invention comprises another inhibitor of HCV polymerase as an other anti-HCV agent.

According to another more specific embodiment, the pharmaceutical composition of this invention comprises an inhibitor of HCV NS3 protease as an other anti-HCV agent.

According to yet another more specific embodiment, the pharmaceutical composition of this invention comprises an inhibitor of another target in the HCV life cycle as an other anti-HCV agent. Examples of such other targets are HCV helicase, HCV NS2/3 protease or HCV IRES.

In an eighth aspect of the invention, there is provided a use of a compound of formula I according to this invention, or of a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment and/or the prevention of a Flaviviridae viral infection, preferably an HCV infection.

25

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The following definitions apply unless otherwise noted:

As used herein, the term (C_{1-n}) alkyl or C_{1-n} alkyl, wherein n is an integer, either alone or in combination with another radical, are intended to mean acyclic straight or branched chain alkyl radicals containing 1 to n carbon atoms respectively. Examples of such radicals include methyl, ethyl, n-propyl, 1-methylethyl (i-propyl), n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl (tert.-butyl), n-pentyl, etc.. In the following the term Me denotes a methyl group.

If an alkyl group is substituted by halogen, it is preferably mono-, di- or trisubstituted with fluorine or monosubstituted by chlorine or bromine. Preferred alkyl-groups which are trisubstituted with fluorine have a terminal CF₃ group.

5

As used herein, the term (C_{2-n}) alkenyl, wherein n is an integer, either alone or in combination with another radical, is intended to mean an unsaturated, acyclic straight chain radical containing two to n carbon atoms, at least two of which are bonded to each other by a double bond. Examples of such radicals are ethenyl (vinyl), 1-propenyl, 2-propenyl, 1-butenyl, etc..

10

As used herein, the term (C_{2-n}) alkynyl, wherein n is an integer, either alone or in combination with another group, is intended to mean an unsaturated, acyclic straight chain alkynyl radical containing 2 to n carbon atoms, at least two of which are bonded to each other by a triple bond. Examples of such radicals are ethynyl, 1-propynyl, 2-propynyl, etc..

15

As used herein, the term (C_{3-n})cycloalkyl, wherein n is an integer, either alone or in combination with another radical, means a cycloalkyl radical containing from three to n carbon atoms and includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

20

As used herein, the term (C_{5-n})cycloalkenyl, wherein n is an integer, either alone or in combination with another radical, means an unsaturated cyclic radical containing five to n carbon atoms. Examples are cyclopentenyl and cyclohexenyl.

25

As used herein the term (C_{1-n})alkyl-(C_{3-m})cycloalkyl, wherein n and m are integers, either alone or in combination with another radical, means a branched or straight chain alkyl radical having 1 to n C-atoms to which a cycloalkyl radical containing from three to m C-atoms is covalently bonded. Preferably the alkyl radical is a straight chain and the cycloalkyl radical is linked to its terminal C-atom. Examples of (C₁₋₃)alkyl-(C₃₋₆)cycloalkyl include cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, 2-cyclopropylethyl, 2-cyclobutylethyl, 2-cyclopentylethyl, 2-cyclohexylethyl, 3-cyclopropylpropyl, 3-cyclobutylpropyl, 3-cyclopentylpropyl, 3-cyclohexylpropyl, etc..

30

35

As used herein, the terms alkyl-aryl, alkyl-HCy, alkyl-Hetaryl, alkyl-Het, etc. mean an alkyl radical to which an aryl, HCy, Hetaryl, Het group is bonded, respectively.

Examples of (C₁₋₃)alkyl-aryl are benzyl (phenylmethyl), phenylethyl and phenylpropyl.

5

As used herein, the term "carboxy protecting group" (CPG) defines protecting groups that can be used during synthetic transformation and are listed in Greene,

"Protective Groups in Organic Chemistry", John Wiley & Sons, New York (1981) and

"The Peptides: Analysis, Synthesis, Biology", Vol. 3, Academic Press, New York


10 (1981), the disclosures of which are hereby incorporated by reference.

A carboxyl group is usually protected as an ester that can be cleaved to give the carboxylic acid. Protecting groups that can be used include: 1) alkyl esters such as methyl, trimethylsilylethyl and *t*-butyl, 2) aralkyl esters such as benzyl and substituted
15 benzyl, or 3) esters that can be cleaved by mild base treatment or mild reductive means such as trichloroethyl and phenacyl esters.


As used herein, the term "aryl" either alone or in combination with another radical means a 6- or 10-membered aryl, i.e. an aromatic radical containing six or ten
20 carbon atoms, for example phenyl, 1-naphthyl or 2-naphthyl. The most preferred meaning of aryl is phenyl.

As used herein the term heteroatom means O, S or N.

25 As used herein, the term "heterocycle", either alone or in combination with another radical, means a monovalent radical derived by removal of a hydrogen from a five-, six-, or seven-membered saturated or unsaturated (including aromatic) heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur. Examples of such heterocycles include, but are not limited to, azetidine, pyrrolidine,
30 tetrahydrofuran, thiazolidine, pyrrole, thiophene, hydantoin, diazepine, 1H-imidazole, isoxazole, thiazole, tetrazole, piperidine, piperazine, homopiperidine, homo-piperazine, 1,4-dioxane, 4-morpholine, 4-thiomorpholine, pyridine, pyridine-N-oxide or pyrimidine, or the following heterocycles:



- As used herein, the term "9- or 10-membered heterobicyclic" or "heterobicyclic" either alone or in combination with another radical, means a heterocycle as defined above fused to one or more other cycle, be it a heterocycle or any other cycle. Examples of such heterobicyclics include, but are not limited to, indole, benzimidazole, thiazolo[4,5-b]-pyridine, quinoline, or coumarin, or the following:



- 10 As used herein, the term "**Het**" defines a 4-, 5-, 6- or 7-membered heterocycle having 1 to 4 heteroatoms selected from O, N and S, which may be saturated, unsaturated or aromatic, or a 8-, 9-, 10- or 11-membered heterobicyclic having 1 to 5 heteroatoms wherever possible, selected from O, N and S, which may be saturated, unsaturated or aromatic, unless specified otherwise.

15

As used herein, the term "**HCy**" defines a saturated or unsaturated 4-, 5-, 6- or 7-membered monocyclic heterocycle having 1 to 3 heteroatoms selected from O, N and S, unless specified otherwise.

- 20 As used herein, the term "**Hetaryl**" defines an aromatic 5- or 6-membered monocyclic heterocycle having 1 or 2 heteroatoms selected from O, N and S, or a 9- or 10-membered aromatic heterobicyclic having 1 to 4 heteroatoms selected from O, N, and S, unless specified otherwise.

- 25 As used herein, the term "halo" means a halogen atom and includes fluorine, chlorine, bromine and iodine.

- As used herein, the term "OH" refers to a hydroxyl group. It is well known to one skilled in the art that hydroxyl groups may be substituted by functional group equivalents. Examples of such functional group equivalents that are contemplated

by this invention include, but are not limited to, ethers, sulfhydryls, and primary, secondary or tertiary amines.

As used herein, the term "SH" refers to a sulfhydryl group. It is intended within the scope of the present invention that, whenever a "SH" or "SR" group is present, it can also be substituted by any other appropriate oxidation state such as SOR, SO₂R, or SO₃R.

It is intended that the term "substituted" when applied in conjunction with a radical having more than one moiety such as C₁₋₆alkyl-aryl, or C₁₋₆alkyl-Het, such substitution applies to both moieties i.e. both the alkyl and aryl or Het moieties can be substituted with the defined substituents.

As used herein, the term "COOH" refers to a carboxylic acid group. It is well known to one skilled in the art that carboxylic acid groups may be substituted by functional group equivalents. Examples of such functional group equivalents that are contemplated by this invention include, but are not limited to, esters, amides, imides, boronic acids, tetrazole, triazoles, N-acylsulfonyldiamides (RCONHSO₂NR₂), and acylsulfonamides (RCONHSO₂R).

As used herein, the term "functional group equivalent" is intended to mean an element or a substituted derivative thereof, that is replaceable by another element that has similar electronic, hybridization or bonding properties.

As used herein, the term "metal catalyst" is intended to mean a metal such as palladium (0) or palladium (2) for use in a cross-coupling reaction. Examples of such palladium catalysts include, but are not limited to, Pd(PPh₃)₄, Pd/C, Pd(OAc)₂, PdCl₂, and the like. Alternative metals that can catalyze cross-coupling reactions include, but are not limited to complexes of Ni, Rh, Ru and Ir, like for example: Ni(acac)₂, Ni(OAc)₂, or NiCl₂.

The term "detectable label" refers to any group that may be linked to the polymerase or to a compound of the present invention such that when the compound is associated with the polymerase target, such label allows recognition either directly or indirectly of the compound such that it can be detected, measured and quantified.

Examples of such "labels" are intended to include, but are not limited to, fluorescent labels, chemiluminescent labels, colorimetric labels, enzymatic markers, radioactive isotopes and affinity tags such as biotin. Such labels are attached to the compound or to the polymerase by well known methods.

5

The term "affinity tag" means a ligand (that is linked to the polymerase or to a compound of the present invention) whose strong affinity for a receptor can be used to extract from a solution the entity to which the ligand is attached. Examples of such ligands include biotin or a derivative thereof, a histidine polypeptide, a polyarginine, an amylose sugar moiety or a defined epitope recognizable by a specific antibody. Such affinity tags are attached to the compound or to the polymerase by well-known methods.

10
The term "photoreactive group" means a group that is transformed, upon activation by light, from an inert group to a reactive species, such as a free radical. Examples of such groups include, but are not limited to, benzophenones, azides, and the like.

15
The term "salt thereof" means any acid and/or base addition salt of a compound according to the invention; preferably a pharmaceutically acceptable salt thereof.

20

The term "pharmaceutically acceptable salt" means a salt of a compound of formula (I) which is, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, generally water or oil-soluble or dispersible, and effective for their intended use. The term includes pharmaceutically-acceptable acid addition salts and pharmaceutically-acceptable base addition salts. Lists of suitable salts are found in, e.g., S.M. Birge et al., J. Pharm. Sci., 1977, 66, pp. 1-19, which is hereby incorporated by reference in its entirety.

25
30

The term "pharmaceutically-acceptable acid addition salt" means those salts which retain the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acid, nitric acid, phosphoric acid, and the like, and organic acids such as acetic acid, trifluoroacetic

35

acid, adipic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, butyric acid, camphoric acid, camphorsulfonic acid, cinnamic acid, citric acid, digluconic acid, ethanesulfonic acid, glutamic acid, glycolic acid, glycerophosphoric acid, hemisulfic acid, hexanoic acid, formic acid, fumaric acid, 2-hydroxyethane-
5 sulfonic acid (isethionic acid), lactic acid, hydroxymaleic acid, malic acid, malonic acid, mandelic acid, mesitylenesulfonic acid, methanesulfonic acid, naphthalene-sulfonic acid, nicotinic acid, 2-naphthalenesulfonic acid, oxalic acid, pamoic acid, pectinic acid, phenylacetic acid, 3-phenylpropionic acid, pivalic acid, propionic acid, pyruvic acid, salicylic acid, stearic acid, succinic acid, sulfanilic acid, tartaric acid, p-
10 toluenesulfonic acid, undecanoic acid, and the like.

The term "pharmaceutically-acceptable base addition salt" means those salts which retain the biological effectiveness and properties of the free acids and which are not biologically or otherwise undesirable, formed with inorganic bases such as ammonia
15 or hydroxide, carbonate, or bicarbonate of ammonium or a metal cation such as sodium, potassium, lithium, calcium, magnesium, iron, zinc, copper, manganese, aluminum, and the like. Particularly preferred are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically-acceptable organic nontoxic bases include salts of primary, secondary, and tertiary
20 amines, quaternary amine compounds, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion-exchange resins, such as methylamine, dimethylamine, trimethylamine, ethylamine, diethylamine, triethylamine, isopropylamine, tripropylamine, tributylamine, ethanolamine, diethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine,
25 lysine, arginine, histidine, caffeine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, tetramethylammonium compounds, tetraethylammonium compounds, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, dibenzylamine, N,N-dibenzylphenethylamine, 1-ephedrine,
30 N,N'-dibenzylethylenediamine, polyamine resins, and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine.

The term "antiviral agent" as used herein means an agent (compound or biological) that
35 is effective to inhibit the formation and/or replication of a virus in a mammal. This

includes agents that interfere with either host or viral mechanisms necessary for the formation and/or replication of a virus in a mammal. Antiviral agents include, for example, ribavirin, amantadine, VX-497 (merimepodib, Vertex Pharmaceuticals), VX-498 (Vertex Pharmaceuticals), Levovirin, Viramidine, Ceplene (maxamine), XTL-001
5 and XTL-002 (XTL Biopharmaceuticals).

The term "other anti-HCV agent" as used herein means those agents that are effective for diminishing or preventing the progression of hepatitis C related symptoms of disease. Such agents can be selected from: immunomodulatory
10 agents, inhibitors of HCV NS3 protease, other inhibitors of HCV polymerase or inhibitors of another target in the HCV life cycle.

The term "immunomodulatory agent" as used herein means those agents (compounds or biologicals) that are effective to enhance or potentiate the immune system response
15 in a mammal. Immunomodulatory agents include, for example, class I interferons (such as α -, β -, δ - and omega interferons, tau-interferons, consensus interferons and asialo-interferons), class II interferons (such as γ -interferons) and pegylated interferons.

The term "inhibitor of HCV NS3 protease" as used herein means an agent (compound
20 or biological) that is effective to inhibit the function of HCV NS3 protease in a mammal. Inhibitors of HCV NS3 protease include, for example, those compounds described in WO 99/07733, WO 99/07734, WO 00/09558, WO 00/09543, WO 00/59929 or WO 02/060926, the Boehringer Ingelheim clinical candidate identified as BILN 2061 and the Vertex pre-development candidate identified as VX-950. Particularly, compounds # 2, 3,
25 5, 6, 8, 10, 11, 18, 19, 29, 30, 31, 32, 33, 37, 38, 55, 59, 71, 91, 103, 104, 105, 112, 113, 114, 115, 116, 120, 122, 123, 124, 125, 126 and 127 disclosed in the table of pages 224-226 in WO 02/060926, can be used in combination with the compounds of the present invention.

30 The term "other inhibitor of HCV polymerase" as used herein means an agent (compound or biological) that is effective to inhibit the function of HCV polymerase in a mammal, whereby this agent has a structure different from the compounds according to this invention and preferably binds to a site of the HCV polymerase different from the site targeted by the compounds according to this invention. Other
35 inhibitors of HCV polymerase include non-nucleosides, for example, those

compounds described in : WO 03/040112 (Rigel), WO 02/100846 A1 (Shire), WO 02/100851 A2 (Shire), WO 01/85172 A1 (GSK), WO 02/098424 A1 (GSK), WO 00/06529 (Merck), WO 02/06246 A1 (Merck), EP 1 256 628 A2 (Agouron).

Furthermore other inhibitors of HCV polymerase also include nucleoside analogs, for
5 example, those compounds described in: WO 01/90121 A2 (Idenix), WO 02/069903 A2 (Biocryst Pharmaceuticals Inc.), WO 02/057287 A2 (Merck/Isis) and WO 02/057425 A2 (Merck/Isis).

The term "inhibitor of another target in the HCV life cycle" as used herein means an
10 agent (compound or biological) that is effective to inhibit the formation and/or replication of HCV in a mammal other than by inhibiting the RNA dependent RNA polymerase of HCV. This includes agents that interfere with either host or HCV viral mechanisms necessary for the formation and/or replication of HCV in a mammal. Inhibitors of another target in the HCV life cycle include, for example, agents that
15 inhibit a target selected from a HCV helicase, HCV NS2/3 protease and HCV IRES. Specific examples of inhibitors of another target in the HCV life cycle include ISIS-14803 (ISIS Pharmaceuticals).

The term "HIV inhibitor" as used herein means an agents (compound or biological) that
20 is effective to inhibit the formation and/or replication of HIV in a mammal. This includes agents that interfere with either host or viral mechanisms necessary for the formation and/or replication of HIV in a mammal. HIV inhibitors include, for example, nucleosidic inhibitors, non-nucleosidic inhibitors, protease inhibitors, fusion inhibitors and integrase inhibitors.

25 The term "HAV inhibitor" as used herein means an agent (compound or biological) that is effective to inhibit the formation and/or replication of HAV in a mammal. This includes agents that interfere with either host or viral mechanisms necessary for the formation and/or replication of HAV in a mammal. HAV inhibitors include Hepatitis A vaccines, for
30 example, Havrix® (GlaxoSmithKline), VAQTA® (Merck) and Avaxim® (Aventis Pasteur).

The term "HBV inhibitor" as used herein means an agent (compound or biological) that is effective to inhibit the formation and/or replication of HBV in a mammal. This includes agents that interfere with either host or viral mechanisms necessary for the formation
35 and/or replication of HBV in a mammal. HBV inhibitors include, for example, agents that

inhibit HBV viral DNA polymerase or HBV vaccines. Specific examples of HBV inhibitors include Lamivudine (Epivir-HBV®), Adefovir Dipivoxil, Entecavir, FTC (Coviracil®), DAPD (DXG), L-FMAU (Clevudine®), AM365 (Amrad), Ldt (Telbivudine), monoval-LdC (Valtorcitabine), ACH-126,443 (L-Fd4C) (Achillion), MCC478 (Eli Lilly),
5 Racivir (RCV), Fluoro-L and D nucleosides, Robustaflavone, ICN 2001-3 (ICN), Bam 205 (Novelos), XTL-001 (XTL), Imino-Sugars (Nonyl-DNJ) (Synergy), HepBzyme; and immunomodulator products such as: interferon alpha 2b, HE2000 (Hollis-Eden), Theradigm (Epimmune), EHT899 (Enzo Biochem), Thymosin alpha-1 (Zadaxin®), HBV DNA vaccine (PowderJect), HBV DNA vaccine (Jefferson Center), HBV antigen
10 (OraGen), BayHep B® (Bayer), Nabi-HB® (Nabi) and Anti-hepatitis B (Cangene); and HBV vaccine products such as the following: Engerix B, Recombivax HB, GenHevac B, Hepacare, Bio-Hep B, TwinRix, Comvax, Hexavac.

The term "class I interferon" as used herein means an interferon selected from a
15 group of interferons that all bind to receptor type I. This includes both naturally and synthetically produced class I interferons. Examples of class I interferons include α -, β -, δ -, omega interferons, tau-interferons, consensus interferons, asialo-interferons.

The term "class II interferon" as used herein means an interferon selected from a group
20 of interferons that all bind to receptor type II. Examples of class II interferons include γ -interferons.

Specific preferred examples of some of these agents are listed below:

- antiviral agents: ribavirin and amantadine;
- 25 ▪ immunomodulatory agents: class I interferons, class II interferons and pegylated interferons;
- HCV NS3 protease inhibitors;
- other inhibitors of the HCV polymerase: nucleosidic and non-nucleosidic inhibitors;
- 30 ▪ inhibitor of another target in the HCV life cycle that inhibits a target selected from: NS3 helicase, HCV NS2/3 protease or internal ribosome entry site (IRES);
- HIV inhibitors: nucleosidic inhibitors, non-nucleosidic inhibitors, protease inhibitors, fusion inhibitors and integrase inhibitors; or
- HBV inhibitors: agents that inhibit viral DNA polymerase or is an HBV vaccine.


As discussed above, combination therapy is contemplated wherein a compound of formula (I), or a pharmaceutically acceptable salt thereof, is co-administered with at least one additional agent selected from: an antiviral agent, an immunomodulatory agent, an inhibitor of HCV NS3 protease, another inhibitor of HCV polymerase, an inhibitor of another target in the HCV life cycle, an HIV inhibitor, an HAV inhibitor and an HBV inhibitor. Examples of such agents are provided in the Definitions section above. These additional agents may be combined with the compounds of this invention to create a single pharmaceutical dosage form. Alternatively these additional agents may be separately administered to the patient as part of a multiple dosage form, for example, using a kit. Such additional agents may be administered to the patient prior to, concurrently with, or following the administration of wherein a compound of formula (I), or a pharmaceutically acceptable salt thereof.

As used herein, the term "treatment" means the administration of a compound or composition according to the present invention to alleviate or eliminate symptoms of the hepatitis C disease and/or to reduce viral load in a patient.

As used herein, the term "prevention" means the administration of a compound or composition according to the present invention post-exposure of the individual to the virus but before the appearance of symptoms of the disease, and/or prior to the detection of the virus in the blood.

The following signs --- and ~ are used interchangeably in subformulas to indicate the bond, or in the case of a spirocyclic group the atom, which is bonded to the rest of the molecule as defined.

As used herein, the designation whereby a bond to a substituent R is drawn as emanating from the center of a ring, such as, for example,




means that the substituent R may be attached to any free position on the ring that would otherwise be substituted by a hydrogen atom, unless specified otherwise.

Preferred embodiments

As long as not stated otherwise, all groups, substituents and indices, like e.g. R^1 , R^2 , R^{2h} , R^3 , R^{4a} , R^{4b} , R^5 , R^{11} , R^{12} , R^{13} , R^{21} , R^{31} , R^{32} , R^{33} , R^{35} , R^{60} , R^{111} , R^{112} , R^{117} , R^{150} , R^{160} , R^{161} , R^{162} , R^{163} , R^O , R^C , R^{N1} , R^{N2} , R^{N3} , R^{N4} , **A**, **B**, M^1 , M^2 , M^3 , M^4 , Y^1 , **Z**, **HCy**, and **Het**, have the definitions as defined hereinbefore and hereinafter. In the following, the preferred embodiments, groups, substituents and indices according to this invention are described.

- 10 In a preferred embodiment of the first aspect of the invention, there is provided an isomer, enantiomer, diastereoisomer or tautomer of a compound, represented by formula I:



15 wherein

----- represents either a single or a double bond;

- 20 **B** is -N- and **A** is CR^1 or =N-; or

B is =C- and **A** is O, S or NR^1 ;

- 25 R^1 is selected from the group consisting of: H, (C_{1-6}) alkyl optionally substituted with:
halogen, OR^{11} , SR^{11} or $N(R^{12})_2$, wherein R^{11} and each R^{12} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**, said aryl or **Het** optionally substituted with R^{160} ; or
both R^{12} are covalently bonded together and to the nitrogen to which they are
30 both attached to form a 5, 6 or 7-membered saturated heterocycle;

the group $-C(=Y^1)-Z$ is covalently linked to either M^2 or M^3 ,

M^1 is CR^{4a} ,

one of M^2 and M^3 is CR^5 ,

5 M^4 is CR^{4b} ,

and in addition one or two of the groups selected from M^1 , M^2 , M^3 and M^4 may also be N, with the proviso that the group M^2 or M^3 to which $-C(=Y^1)-Z$ is linked is an C-atom,

10

Y^1 is O or S;

Z is defined as $NR^{N2}-SO_2-R^C$, wherein R^C is optionally substituted with R^{60} ;

15 R^2 is selected from: halogen or R^{21} , wherein R^{21} is aryl or **Het**, said R^{21} is optionally substituted with R^{150} ;

R^3 is selected from (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl, (C_{5-7}) cycloalkenyl, (C_{1-3}) alkyl- (C_{5-7}) cycloalkenyl, (C_{6-10}) bicycloalkyl, (C_{1-3}) alkyl- (C_{6-10}) bicycloalkyl, (C_{6-10}) bicycloalkenyl, (C_{1-3}) alkyl- (C_{6-10}) bicycloalkenyl, **HCy** or (C_{1-3}) alkyl-**HCy**,

20

wherein **HCy** is a saturated or unsaturated 4 to 7-membered heterocyclic group with 1 to 3 heteroatoms selected from O, S and N;

said alkyl, cycloalkyl, cycloalkenyl, bicycloalkyl, bicycloalkenyl, **HCy** and

25

alkyl-**HCy** being optionally substituted with from 1 to 4 substituents selected from: a) halogen;

b) (C_{1-6}) alkyl optionally substituted with:

- OR^{31} or SR^{31} wherein R^{31} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl; or

30

- $N(R^{32})_2$ wherein each R^{32} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl; or both R^{32} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

- c) OR^{33} or SR^{33} wherein R^{33} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl;
- d) $\text{N}(\text{R}^{35})_2$ wherein each R^{35} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl; or both R^{35} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

R^{4a} , R^{4b} , R^5 each are independently H or defined as R^{150} ;

- 10 R^{60} is defined as 1 to 4 substituents independently selected from:
- 1 to 3 substituents selected from halogen;
 - one of each substituent selected from: OPO_3H , NO_2 , cyano, azido, $\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6})$ alkyl or $\text{C}(=\text{NH})\text{NHCO}(\text{C}_{1-6})$ alkyl, SO_3H ; and
 - 1 to 3 substituents selected from:
- 15 a) (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, C_{3-7} spirocycloalkyl optionally containing 1 or 2 heteroatom selected from N, O and S; (C_{2-6}) alkenyl, (C_{2-8}) alkynyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, all of which optionally being substituted with R^{150} ;
- b) OR^0 ;
- 20 c) $\text{OC}(\text{O})\text{R}^0$;
- d) SR^0 , SO_2R^C , $\text{SO}_2\text{N}(\text{R}^{N2})\text{R}^{N1}$, $\text{SO}_2\text{N}(\text{R}^{N2})\text{C}(\text{O})\text{R}^C$ or $\text{CONR}^{N2}\text{SO}_2\text{R}^C$;
- e) $\text{N}(\text{R}^{N2})\text{R}^{N1}$, $\text{N}(\text{R}^{N2})\text{COOR}^C$ or $\text{N}(\text{R}^{N2})\text{SO}_2\text{R}^C$;
- f) $\text{N}(\text{R}^{N2})\text{COR}^C$;
- g) $\text{N}(\text{R}^{N3})\text{CON}(\text{R}^{N2})\text{R}^{N1}$;
- 25 h) $\text{N}(\text{R}^{N3})\text{COCOR}^C$, $\text{N}(\text{R}^{N3})\text{COCOOR}^0$ or $\text{N}(\text{R}^{N3})\text{COCON}(\text{R}^{N2})\text{R}^{N1}$;
- i) COR^0 ;
- j) COOR^0 ;
- k) $\text{CON}(\text{R}^{N2})\text{R}^{N1}$;
- 30 l) aryl, Het, (C_{1-4}) alkyl)aryl or (C_{1-4}) alkyl)Het, all of which optionally being substituted with R^{150} ;
- wherein said R^{N1} , R^C and/or R^0 are optionally substituted with R^{150} as defined,

R^{150} is defined as 1 to 4 substituents independently selected from:

- 1 to 3 substituents selected from halogen;
 - one of each substituent selected from: OPO_3H , NO_2 , cyano, azido, $\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6})\text{alkyl}$ or $\text{C}(=\text{NH})\text{NHCO}(\text{C}_{1-6})\text{alkyl}$; and
 - 1 to 3 substituents selected from:
- 5 a) $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $\text{C}_{3-7}\text{spirocycloalkyl}$ optionally containing 1 or 2 heteroatoms selected from N, O and S; $(\text{C}_{2-6})\text{alkenyl}$, $(\text{C}_{2-8})\text{alkynyl}$, $(\text{C}_{1-3})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, all of which optionally substituted with R^{160} ;
- b) OR^{O} ;
- c) $\text{OC}(\text{O})\text{R}^{\text{O}}$;
- 10 d) SR^{O} , $\text{SO}_2\text{R}^{\text{C}}$, $\text{SO}_2\text{N}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$ or $\text{SO}_2\text{N}(\text{R}^{\text{N}2})\text{C}(\text{O})\text{R}^{\text{C}}$;
- e) $\text{N}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$, $\text{N}(\text{R}^{\text{N}2})\text{COOR}^{\text{C}}$ or $\text{N}(\text{R}^{\text{N}2})\text{SO}_2\text{R}^{\text{C}}$;
- f) $\text{N}(\text{R}^{\text{N}2})\text{COR}^{\text{C}}$;
- g) $\text{N}(\text{R}^{\text{N}3})\text{CON}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$;
- h) $\text{N}(\text{R}^{\text{N}3})\text{COCOR}^{\text{C}}$, $\text{N}(\text{R}^{\text{N}3})\text{COCOOR}^{\text{O}}$ or $\text{N}(\text{R}^{\text{N}3})\text{COCON}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$;
- 15 wherein $\text{R}^{\text{N}1}$ is as defined or OH, OAlkyl;
- i) COR^{O} ;
- j) COOR^{O} ;
- k) tetrazole or $\text{CON}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$;
- wherein said $\text{R}^{\text{N}1}$, R^{C} and/or R^{O} are optionally substituted with R^{160} as
- 20 defined;

R^{160} is defined as 1, 2 or 3 substituents independently selected from:

- 1, 2 or 3 fluorine substituents; and
 - one of each substituent selected from tetrazole, chlorine, bromine, iodine,
- 25 CN, nitro, $\text{C}_{1-4}\text{alkyl}$, CF_3 , COOR^{161} , SO_3H , SR^{161} , $\text{SO}_2\text{R}^{163}$, OR^{161} , $\text{N}(\text{R}^{162})_2$, $\text{SO}_2\text{N}(\text{R}^{162})_2$, $\text{SO}_2\text{NR}^{162}\text{COR}^{162}$, $\text{NR}^{162}\text{SO}_2\text{R}^{163}$, $\text{NR}^{162}\text{COR}^{162}$ or $\text{CON}(\text{R}^{162})_2$, wherein R^{161} , R^{163} and each R^{162} is independently $(\text{C}_{1-4})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-3})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$; and R^{161} and each R^{162} may each independently also be H; or both R^{162} are covalently bonded together and to
- 30 the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

R^{O} , R^{C} are independently defined as $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-6})\text{cycloalkyl}$, $(\text{C}_{1-4})\text{alkyl}-(\text{C}_{3-6})\text{cycloalkyl}$, $(\text{C}_{2-6})\text{alkenyl}$, aryl, Het, $(\text{C}_{1-4})\text{alkyl-aryl}$, $(\text{C}_{1-4})\text{alkyl-Het}$;

R^{N1} is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₄)alkyl-(C₃₋₆)cycloalkyl, (C₂₋₆)alkenyl, aryl, Het, (C₁₋₄)alkyl-aryl, (C₁₋₄)alkyl-Het; or

- 5 R^{N2} , R^{N3} , R^{N4} are independently H, CH₃, (C₂₋₆)alkyl, (C₃₋₆)cycloalkyl, (C₁₋₄)alkyl-(C₃₋₆)cycloalkyl; all of which being optionally substituted with halogen, carboxy or C₁₋₆-alkoxycarbonyl; and/or wherein said alkyl, cycloalkyl or alkylcycloalkyl is optionally substituted with hydroxy, C₁₋₆-alkyl, C₁₋₆-alkoxy, amino, -NH(C₁₋₄-alkyl) and/or -N(C₁₋₄-alkyl)₂; and

10

in the case

- a) of a group N(R^{N2})(R^{N1}) the substituents R^{N2} and R^{N1} ; or
 b) of a group NR^{N3}-N(R^{N2})(R^{N1}) the substituents R^{N3} and R^{N1} , or R^{N2} and R^{N1} ; may be covalently bonded together to form a 4-, 5-, 6- or 7-membered
 15 saturated or unsaturated N-containing heterocycle or a 8-, 9-, 10- or 11-membered N-containing heterobicyclic each may have additionally from 1 to 3 heteroatoms selected from O, N, and S, wherein said heterocycle or heterobicyclic is optionally substituted as defined;


- 20 wherein Het is defined as a 4-, 5-, 6- or 7-membered heterocycle having 1 to 4 heteroatoms selected from O, N and S, or a 8-, 9-, 10- or 11-membered heterobicyclic having 1 to 5 heteroatoms selected from O, N and S;

or a salt thereof.


25

Core:






This invention comprises compounds of the formulas Ia and Ib



30



Furthermore this invention comprises compounds according to the formulas I.1 to I.5













Preferably the groups M^1 and M^4 are CR^{4a} and CR^{4b} , respectively.

One group of M^2 and M^3 to which the group $-C(=Y^1)-Z$ is covalently linked is preferably C and the other group of M^3 and M^2 is preferably CR^5 .

5






Therefore those compounds are preferred which are described by the following formulas





whereby the fused phenyl-ring may be mono-, di- or tri-substituted, preferably mono- or di-substituted, with R^{150} .











- 5 The most preferred compounds of the beforementioned formulas are described by the following formulas:





- 10 whereby the fused phenyl ring may be mono-, di- or tri-substituted, preferably mono- or di-substituted, with R^{150} .

Furthermore compounds defined by the following formulas also belong to this

invention



Yet furthermore, compounds defined by the following formulas also belong to this invention:



R¹:

According to a preferred embodiment of this invention **R¹** is selected from the group consisting of H and (C₁₋₆)alkyl, especially of H, CH₃, ethyl and isobutyl. Most preferably **R¹** is H or methyl, in particular methyl.

5

Y¹:

A preferred definition of **Y¹** is O.

Z:

- 10 According to a preferred embodiment **Z** is defined as **NR^{N3}-SO₂-N(R^{N2})R^{N1}**, wherein **R^{N1}**, **R^{N2}** and **R^{N3}** are defined as hereinbefore and **R^{N1}** or any heterocycle formed by **R^{N1}** and **R^{N2}** is optionally substituted with **R⁶⁰**.

- According to a second preferred embodiment **Z** is defined as **NR^{N2}-SO₂-R^C**, wherein
15 **R^C** and **R^{N2}** are defined as hereinbefore and **R^C** is optionally substituted with **R⁶⁰**.
Most preferably, **Z** is defined as **NR^{N2}-SO₂-R^C** wherein **R^{N2}** and **R^C** are defined as follows:

- R^{N2}** is H, (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl or (C₁₋₃)alkyl-(C₃₋₆)cycloalkyl; in particular H or
20 methyl; most preferably H; and

- R^C** is (C₁₋₆)alkyl, (C₃₋₆)cycloalkyl, (C₁₋₃)alkyl-(C₃₋₆)cycloalkyl, (C₂₋₆)alkenyl, phenyl,
naphthyl, **Het**, (C₁₋₃)alkyl-phenyl, (C₁₋₃)alkyl-naphthyl, (C₁₋₃)alkyl-**Het**, wherein
25 said alkyl, cycloalkyl, alkyl-cycloalkyl, alkenyl, phenyl, naphthyl, **Het**, alkyl-phenyl, alkyl-naphthyl, or alkyl-**Het**, are all optionally substituted with 1 to 4
substituents selected from **R⁶⁰**; preferably selected from **R¹⁵⁰**.

- In this embodiment the preferred definition of **R^C** is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopropylmethyl, cyclobutylmethyl,
30 cyclopentylmethyl, pyrrolidine, piperidine, morpholine, thiomorpholine, piperazine, phenyl, naphthyl, benzyl, thiophene, furan, pyrrole, imidazole, pyrazole, oxazole, isoxazole, thiazole, pyridazine, pyrimidine, pyrazine, diazepine, azepine, quinoline, isoquinoline, benzofuran, benzothiophene, benzothiazole, purine, pteridine,

2,1,3-benzothiadiazole  , and


imidazo[2,1-B][1,3]thiazole  ;


all of which are optionally substituted with 1 to 3 substituents selected from R^{60} ; preferably selected from R^{150} . Most preferred substituents thereof are halogen, nitro, (C₁₋₃)alkyl, O(C₁₋₃)alkyl, carboxyl, COO(C₁₋₃)alkyl, NHCO(C₁₋₃)alkyl, wherein the alkyl groups may be substituted by halogen.

R²:

Preferably R^2 is R^{21} , wherein R^{21} is a phenyl or Het selected from the group of formulas

10






and wherein said R^{21} is unsubstituted or substituted with R^{150} .

5

A very most preferred definition of R^2 is



defined.

In the case R^2 as defined above is substituted, it is preferably substituted with 1, 2 or 3 substituents selected from:

- 10 - 1 to 3 substituents selected from halogen;
- one of each substituent selected from: NO_2 , cyano, azido; and
- 1 to 2 substituents selected from:
- a) (C_{1-4}) alkyl or (C_{1-4}) alkoxy, both optionally substituted with OH, $O(C_{1-4})$ alkyl, $SO_2(C_{1-4})$ alkyl, 1 to 3 halogen atoms, amino, $NH(CH_3)$ or $N(CH_3)_2$;
- 15 b) $NR^{111}R^{112}$ wherein both R^{111} and R^{112} are independently H, (C_{1-4}) alkyl, or R^{112} is (C_{3-7}) cycloalkyl, (C_{1-3}) alkyl (C_{3-7}) cycloalkyl, phenyl, benzyl; or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle, each of said alkyl, cycloalkyl, alkylcycloalkyl, phenyl and benzyl, being optionally substituted with halogen or:

-OR^{2h} or N(R^{2h})₂, wherein each R^{2h} is independently H, (C₁₋₄)alkyl, or both R^{2h} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle;

- c) NHCOR¹¹⁷ wherein R¹¹⁷ is (C₁₋₄)alkyl, O(C₁₋₄)alkyl or O(C₃₋₇)cycloalkyl; and
 5 e) CONH₂, CONH(C₁₋₄)alkyl, CON((C₁₋₄)alkyl)₂.

Most preferred substituents of R² are selected from:

- 1 to 2 substituents selected from fluorine;
- one of each substituent selected from: chlorine, bromine, NO₂, cyano; and
- 10 - 1 to 2 substituents selected from:
 - a) methyl, trifluoromethyl, ethyl, n-propyl, i-propyl, methoxy, trifluoromethoxy, ethoxy, n-propoxy or i-propoxy, wherein said methyl, ethyl, n-propyl, i-propyl, ethoxy, n-propoxy and i-propoxy are optionally substituted with OH, methoxy, amino, NH(CH₃) or N(CH₃)₂;
 - 15 b) NR¹¹¹R¹¹² wherein both R¹¹¹ and R¹¹² are independently H or methyl, or R¹¹² is phenyl or benzyl;
 - c) NHCOR¹¹⁷ wherein R¹¹⁷ is methyl or methoxy; and
 - e) CONH₂, CONH(CH₃), CON(CH₃)₂.


20

R³:

R³ is preferably selected from (C₃₋₇)cycloalkyl, (C₅₋₇)cycloalkenyl, (C₆₋₁₀)bicycloalkyl, (C₆₋₁₀)bicycloalkenyl, or HCy, wherein said groups are unsubstituted or mono- or disubstituted by halogen, hydroxy, (C₁₋₄)alkyl and/or O-(C₁₋₄)alkyl, wherein the alkyl
 25 groups may be fluorinated.

Most preferably R³ is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl,

or a group selected from:



wherein all said cyclic groups are unsubstituted or substituted by fluorine, (C₁₋₃)alkyl
 30 or CF₃.

The very most preferred definition of R^3 is cyclopentyl or cyclohexyl.

R^{4a} , R^{4b} , R^5 :

- 5 Preferably R^{4a} , R^{4b} , R^5 each are independently H, hydroxy, halogen, cyano, nitro, carboxyl, (C_{1-4}) alkyl, CF_3 , (C_{1-4}) alkoxy, -O- (C_{3-7}) cycloalkyl, -O- (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl, -O-aryl, -O- (C_{1-3}) alkyl-aryl, -O-Het, -O- (C_{1-3}) alkyl-Het, $NR^{N1}R^{N2}$, or COR^O , $NR^{N2}COR^C$, $CONR^{N2}R^{N1}$, $NR^{N3}CONR^{N1}R^{N2}$, in particular $NHCO(C_{1-4})$ alkyl or $CONHR^{N1}$, $NHCONHR^{N1}$;
- 10 wherein R^C , R^O , R^{N1} , R^{N2} , R^{N3} are as defined; preferably R^C , R^O and R^{N1} are independently of each other H, (C_{1-4}) alkyl, aryl, (C_{1-3}) alkyl-aryl, wherein aryl is preferably optionally substituted phenyl; and preferably R^{N2} and R^{N3} are independently H or methyl; wherein all said alkyl groups, including alkoxy, may be mono-, di- or trisubstituted by fluorine or mono-substituted by chlorine or bromine.
- 15 Most preferred substituents R^{4a} , R^{4b} , R^5 each are independently H, hydroxy, halogen, cyano, nitro, methyl, CF_3 , methoxy, carboxy, amino, -NMe₂, -CONH₂, -NHCONH₂, -CO-NHMe, -NHCONHMe, -CO-NMe₂ or -NHCONMe₂; in particular H, methyl or methoxy. Preferably R^{4a} is H or methyl. Very most preferably at least two of the
- 20 substituents selected from R^{4a} , R^{4b} , R^5 are H.

R^{60} :

- The substituents R^{60} are preferably each defined as 1 to 4 substituents
- 25 independently selected from:
- 1 to 3 substituents selected from halogen;
 - one of each substituent selected from: NO₂, cyano, azido; and
 - 1 to 3 substituents selected from:
- a) (C_{1-4}) alkyl, (C_{3-7}) cycloalkyl, (C_{2-4}) alkenyl, (C_{2-4}) alkynyl, (C_{1-3}) alkyl- (C_{3-7}) cycloalkyl,
- 30 all of which optionally being substituted with R^{150} ;
- b) OR^O ;
 - e) $N(R^{N2})R^{N1}$;
 - f) $N(R^{N2})COR^C$;
 - j) $COOR^O$;

k) $\text{CON}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$; .

l) phenyl, Het, $(\text{C}_{1-3}\text{alkyl})\text{phenyl}$ or $(\text{C}_{1-3}\text{alkyl})\text{Het}$; wherein

Het is selected from furan, tetrahydrofuran, thiophene, tetrahydrothiophene, tetrahydropyran, pyridinyl, azetidine, pyrrolidine, piperidine, piperazine, morpholine,
5 thiomorpholine, homopiperidine and homopiperazine;

wherein said $\text{R}^{\text{N}1}$, R^{C} and/or R^{O} are optionally substituted with R^{150} as defined.

R^{150} :

- 10 R^{150} is preferably defined as 1 to 4 substituents independently selected from:
- 1 to 3 fluorine-substituents;
 - one of each substituent selected from: chlorine, bromine, iodine, NO_2 , cyano, azido;
 - and
 - 1 to 3 substituents selected from:
- 15 a) (C_{1-3}) alkyl, CF_3 , $(\text{C}_{3-6})\text{cycloalkyl}$, (C_{1-3}) alkyl- $(\text{C}_{3-6})\text{cycloalkyl}$, all of which optionally substituted with R^{160} ;
- b) OR^{O} ;
- e) $\text{N}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$;
- f) $\text{N}(\text{R}^{\text{N}2})\text{COR}^{\text{C}}$;
- 20 j) COOR^{O} ;
- k) $\text{CON}(\text{R}^{\text{N}2})\text{R}^{\text{N}1}$;

wherein said $\text{R}^{\text{N}1}$, R^{C} and/or R^{O} are optionally substituted with R^{160} as defined.

25 R^{160} :

R^{160} is preferably defined as 1, 2 or 3 substituents independently selected from:

- 1, 2 or 3 fluorine substituents; and
 - one of each substituent selected from chlorine, bromine, iodine, CN, nitro, methyl, trifluoromethyl, ethyl, n-propyl, i-propyl, COOH , COOCH_3 , OH , OCH_3 , OCF_3 , NH_2 ,
- 30 NHCH_3 , $\text{N}(\text{CH}_3)_2$, SO_2NH_2 , NHCOCH_3 , $\text{SO}_2\text{NHCOCH}_3$ or CONH_2 , CONHCH_3 and $\text{CON}(\text{CH}_3)_2$.

R^{O} , R^{C} :

- 35 Preferably R^{O} , R^{C} are independently selected from $(\text{C}_{1-4})\text{alkyl}$, $(\text{C}_{3-6})\text{cycloalkyl}$,

(C₁₋₃)alkyl-(C₃₋₆)cycloalkyl, phenyl, benzyl, **Het**, (C₁₋₃)alkyl-**H t**; all of which are optionally substituted as defined; and R^o may also be H.

5 R^{N1}, R^{N2}, R^{N3}, R^{N4}.

R^{N1} is preferably selected from H, (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl, (C₁₋₃)alkyl-(C₃₋₆)cycloalkyl, phenyl, benzyl, phenylethyl, **Het**, (C₁₋₃)alkyl-**Het**; wherein said alkyl, cycloalkyl, alkyl-cycloalkyl, phenyl, benzyl, phenylethyl, **Het** and alkyl-**Het** are optionally substituted as defined; or

10

R^{N2}, R^{N3}, R^{N4} are independently selected from H, methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopropylmethyl; all of which being optionally substituted with methyl, fluorine, carboxy or methoxycarbonyl; and/or wherein said ethyl, n-propyl or i-propyl, but preferably not the C-atom thereof directly bonded to the

15 N-atom, is optionally substituted with hydroxy, methyl, methoxy, amino, -NH(CH₃) and/or -N(CH₃)₂; and

in the case

- a) of a group N(R^{N2})R^{N1} the substituents R^{N2} and R^{N1}; or
- 20 b) of a group NR^{N3}-N(R^{N2})R^{N1} the substituents R^{N3} and R^{N1} or R^{N2} and R^{N1}; may be covalently bonded together to form a 5-, 6- or 7-membered saturated heterocycle which may have additionally one heteroatom selected from O, N, and S, wherein said heterocycle is optionally substituted as defined.

25 Included within the scope of this invention are all compounds of formula I as presented in Tables 1 and 2. Preferred compounds of these tables show an IC₅₀ value of below 1 μM (range B as defined hereinafter).

30 **Polymerase activity**

The ability of the compounds of formula (I) to inhibit RNA synthesis by the RNA dependent RNA polymerase of HCV can be demonstrated by any assay capable of measuring RNA dependent RNA polymerase activity. A suitable assay is described in the examples.

35

Specificity for RNA dependent RNA polymeras activity

To demonstrate that the compounds of the invention act by specific inhibition of HCV polymerase, the compounds may be tested for inhibitory activity in an assay measuring the activity of an RNA-dependent RNA polymerase other than the HCV
5 polymerase or in a DNA dependent RNA polymerase assay.

When a compound of formula (I), or one of its therapeutically acceptable salts, is employed as an antiviral agent, it is administered orally, topically or systemically to mammals, e.g. humans, cattle, pig, dogs, cats, rabbits or mice, in a vehicle
10 comprising one or more pharmaceutically acceptable carriers, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard biological practice.

For oral administration, the compound or a therapeutically acceptable salt thereof
15 can be formulated in unit dosage forms such as capsules or tablets each containing a predetermined amount of the active ingredient, ranging from about 25 to 500 mg, in a pharmaceutically acceptable carrier.

For topical administration, the compound can be formulated in pharmaceutically
20 accepted vehicles containing 0.1 to 5 percent, preferably 0.5 to 5 percent, of the active agent. Such formulations can be in the form of a solution, cream or lotion.

For parenteral administration, the compound of formula (I) is administered by either intravenous, subcutaneous or intramuscular injection, in compositions with
25 pharmaceutically acceptable vehicles or carriers. For administration by injection, it is preferred to use the compounds in solution in a sterile aqueous vehicle which may also contain other solutes such as buffers or preservatives as well as sufficient quantities of pharmaceutically acceptable salts or of glucose to make the solution isotonic.

30
Suitable vehicles or carriers for the above noted formulations are described in pharmaceutical texts, e.g. in "Remington's The Science and Practice of Pharmacy", 19th ed., Mack Publishing Company, Easton, Penn., 1995, or in "Pharmaceutical Dosage Forms And Drugs Delivery Systems", 6th ed., H.C. Ansel et al., Eds.,
35 Williams & Wilkins, Baltimore, Maryland, 1995.

The dosage of the compound will vary with the form of administration and the particular active agent chosen. Furthermore, it will vary with the particular host under treatment. Generally, treatment is initiated with small increments until the optimum
5 effect under the circumstance is reached. In general, the compound of formula I is most desirably administered at a concentration level that will generally afford antivirally effective results without causing any harmful or deleterious side effects.

For oral administration, the compound or a therapeutically acceptable salt is
10 administered in the range of 10 to 200 mg per kilogram of body weight per day, with a preferred range of 25 to 150 mg per kilogram.

For systemic administration, the compound of formula (I) is administered at a dosage of 10 mg to 150 mg per kilogram of body weight per day, although the
15 aforementioned variations will occur. A dosage level that is in the range of from about 10 mg to 100 mg per kilogram of body weight per day is most desirably employed in order to achieve effective results.

When the compositions of this invention comprise a combination of a compound of
20 formula I and one or more additional therapeutic or prophylactic agent, both the compound and the additional agent should be present at dosage levels of between about 10 to 100%, and more preferably between about 10 and 80% of the dosage normally administered in a monotherapy regimen.

When these compounds or their pharmaceutically acceptable salts are formulated
25 together with a pharmaceutically acceptable carrier, the resulting composition may be administered *in vivo* to mammals, such as man, to inhibit HCV polymerase or to treat or prevent HCV virus infection. Such treatment may also be achieved using the compounds of this invention in combination with agents which include, but are not limited to: immunomodulatory agents, such as α -, β -, or γ -interferons; other antiviral
30 agents such as ribavirin, amantadine; other inhibitors of HCV NS5B polymerase; inhibitors of other targets in the HCV life cycle, which include but are not limited to, helicase, NS2/3 protease, NS3 protease, or internal ribosome entry site (IRES); or combinations thereof. The additional agents may be combined with the compounds of this invention to create a single dosage form. Alternatively these additional agents
35 may be separately administered to a mammal as part of a multiple dosage form.

Meth d logy and Synthesis

The synthesis of compounds according to this invention is preferably accomplished
5 by applying the procedures as outlined in the examples, including any adaptation of
these procedures and/or applying additional synthesis steps known to the person
skilled in the art. For example, carboxylic acid or thiocarboxylic acid intermediates (I"
in the schemes described hereinafter) required in the preparation of compounds of
formula I can be prepared by procedures, or adaptations thereof, described in WO
10 03/010141 or WO 02/04425.

EXAMPLES

The present invention is illustrated in further detail by the following non-limiting
examples. All reactions are performed in a nitrogen or argon atmosphere.
15 Temperatures are given in degrees Celsius. Flash chromatography is performed on
silica gel. Solution percentages or ratios express a volume to volume relationship,
unless stated otherwise. Mass spectral analyses are recorded using electrospray
mass spectrometry. Hereinbefore and hereinafter the following abbreviations or
symbols are used:


- 20 AcOH: acetic acid;
BOP: benzotriazole-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate;
BroP: Bromo tris(dimethylamino)phosphonium hexafluorophosphate;
Bu: butyl;
25 DCC: 1,3-Dicyclohexyl carbodiimide;
DCM: dichloromethane;
DIEA: diisopropylethylamine;
DMAP: 4-(dimethylamino)pyridine;
DMF: *N,N*-dimethylformamide;
30 DMSO: dimethylsulfoxide;
EDAC: see EDC;
EDC: 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride;
ES⁻: electro spray (negative ionization);
ES⁺: electro spray (positive ionization);
35 Et: ethyl;

- Et₂O: diethyl ether;
EtOAc: ethyl acetate;
EtOH: ethanol ;
HATU: O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
5 hexafluorophosphate;
HPLC: high performance liquid chromatography;
ⁱPr: isopropyl ;
ⁱPrOH: isopropanol;
Me: methyl;
10 MeCN: acetonitrile;
MeOH: methanol;
MS (ES): electrospray mass spectrometry;
Ph: phenyl;
Ra-Ni: Raney nickel
15 RT: room temperature (approximatly 25°C)
TBME: *tert*-butylmethyl ether;
TBTU: 2-(1*H*-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate;
tBu: *tert*.-butyl;
TFA: trifluoroacetic acid;
20 THF: tetrahydrofuran;
TLC: thin layer chromatography;

EXAMPLE 1

General procedure for the preparation of acylsulfonamides of general formula I:

- 25 Acylsulfonamide derivatives of the general formula I ($Z = \text{NR}^{\text{N}2}\text{SO}_2\text{R}^{\text{C}}$) according to this invention are preferably prepared by firstly activating the carboxyl group (or thio analog) of the corresponding carboxylic acid derivative (or thio analog) of formula I' and secondly condensing the activated carboxylic acid derivative (or thio analog)
30 with sulfonamides using standard procedures as depicted in the following scheme.



wherein A, B, M¹ to M⁴, R², R³, Y¹, Z, R^{N2} and R^c are as defined hereinbefore, Z is defined as NR^{N2}SO₂R^c and --- represents either a single or double bond.

5

The carboxylic acid derivative (or thio analog) of formula I', including esters thereof, are preferably synthesized according or in analogy to the procedures as outlined in WO 03/010141 and WO 02/04425.


- 10 Activation of carboxylic acids such as those described in the scheme above, can be accomplished using standard amide bond forming reagents such as TBTU, HATU, BOP, BroP, EDAC, DCC, isobutyl chloroformate and the like. Alternatively, activation of the carboxyl group is accomplished by conversion to the corresponding acid chloride prior to condensation with a sulfonamide. Sulfonamides can be
- 15 obtained from commercial sources or prepared from the corresponding sulfonyl chlorides using a solution of ammonia in an inert solvent such as dioxane, THF and the like.

EXAMPLE 2

- 20 General procedure for the preparation of acylsulfonyldiamides of general formula I:

Acylsulfonyldiamide derivatives of general formula I (Z = NR^{N3}SO₂N(R^{N2})R^{N1}) are prepared by condensation of carboxylic acid derivative of general formula I' (or thio analog) with sulfonyldiamides using standard procedures as depicted in the scheme

25 below.




wherein A, B, M^1 to M^4 , R^2 , R^3 , Y^1 , R^{N1} , R^{N2} and R^{N3} are as defined hereinbefore, Z is defined as $NR^{N3}SO_2N(R^{N2})R^{N1}$ and --- represents either a single or double bond.

- 5 The carboxylic acid derivative (or thio analog) of formula I', including esters thereof, are preferably synthesized according or in analogy to the procedures as outlined in WO 03/010141 and WO 02/04425.

- 10 Activation of carboxylic acids such as those described in the scheme, can be accomplished using standard amide bond forming reagents such as TBTU, HATU, BOP, BroP, EDAC, DCC, isobutyl chloroformate and the like. Alternatively, activation of the carboxyl group is accomplished by conversion to the corresponding acid chloride prior to condensation with a sulfonyldiamide. Sulfonyldiamides of formula $HNR^{N3}SO_2N(R^{N2})R^{N1}$, as required in the second step of the scheme, can be prepared by known methods, such as heating amines of formula $R^{N3}NH_2$ with sulfamide in an appropriate solvent (e.g. dioxane) at temperatures ranging from 80 to 160 °C. Such procedures are exemplified in the literature (e.g. R. Sarges *Journal of Medicinal Chemistry* 1976, 19, 695-709).

20 EXAMPLE 3



Methyl 4-amino-3-iodobenzoate :

- Methyl 4-aminobenzoate (30.23 g, 200 mmol) was added to a mixture of iodine (60.91 g, 240 mmol) and silver sulfate (81.07 g, 260 mmol) in 1 L of ethanol at room temperature. The mixture was stirred at room temperature for 18 hours, filtered over
5 Celite and the filtrate was evaporated to dryness under reduced pressure. The residue was chromatographed on silica gel (230-400 mesh) and eluted with 1% ethyl acetate in hexane to give 35 g (63%).

Methyl 4-(N-cyclopentylamino)-3-iodobenzoate:

- 10 A mixture of methyl 4-amino-3-iodobenzoate from above (27.69 g, 100 mmol), cyclopentanone (52.8 mL, 600 mmol), and anhydrous sodium sulfate (1000 g, 142 mmol) in 500 mL of glacial acetic acid was stirred for two hours. Solid sodium triacetoxyborohydride (76.92, 345 mmol) was added in portions and stirred for 16 hrs at room temperature. The mixture was poured on to cool 900 mL of saturated
15 aqueous sodium bicarbonate solution carefully and extracted with 150 mL (3x) of ethyl acetate. The ethyl acetate fraction was dried and evaporated. Purification with silica gel and 0.5% ethyl acetate in hexane gave 24 g (70%).

Methyl 1-cyclopentyl-3-methyl-2-trimethylsilylindole-5-carboxylate:

- 20 A mixture of methyl 4-(N-cyclopentylamino)-3-iodobenzoate from above (22.42 g, 65 mmol), palladium acetate (738 mg, 3.28 mmol), lithium chloride (2.757 g, 65 mmol), potassium acetate (12.757 g, 92.30 mmol), and 1-trimethylsilyl-1-propyne (28.70 mL, 192.56 mmol) in 390 mL of DMF was heated at 100 °C for 6 hours. The mixture was added onto 500 mL of saturated aqueous ammonium chloride solution and extracted
25 with ethyl acetate. The ethyl acetate fraction was dried, evaporated and semi-purified by flash chromatography using silica gel and 0.1% ethyl acetate in hexane. Adding hexane to the semi-purified compound gave 16 g of white crystals (75%).

Methyl 2-bromo-1-cyclopentyl-3-methylindole-5-carboxylate:

- 30 1-Cyclopentyl-3-methyl-2-trimethylsilylindole-5-carboxylate from above (16.45 g, 50 mmol) was dissolved in 500 mL of dichloromethane, cooled to 0 °C. A solution of bromine (1 M, 45 mL, 45 mmol) in dichloromethane was added slowly and stirred for 30 minutes. The solvent was evaporated under reduced pressure, chromatographed on silica gel with 0.1% ethyl acetate to give a semi-pure
35 compound. Addition of hexane gave 12 g (71%).

Methyl 1-cyclopentyl-3-methyl-2-(2-pyridyl)indole-5-carboxylate :


The 2-bromoindole from above (1.53 g, 4.55 mmol, 1.0 equiv.) was dissolved in anhydrous THF (20 mL) and the solution was cooled to -78 °C under an argon atmosphere. *N*-BuLi (solution in hexane, 4.78 mmol, 1.05 equiv.) was added dropwise and the mixture was stirred 30 min in the cold. Trimethylborate (0.62 mL, 5.46 mmol, 1.2 equiv.) was added dropwise and the solution stirred at -78 °C for 40 min and then at 0 °C for 1 h. Finally the reaction mixture was allowed to warm up to room temperature and additional THF (20 mL) was added. *para*-Tolylphosphine (0.083 g, 0.27 mmol, 0.06 equiv.) and 2-bromopyridine (0.86 g, 5.46 mmol, 1.2 equiv.) were then added and the solution was degassed by bubbling argon through for 30 min. Palladium acetate (10 mg, 0.046 mmol, 0.01 equiv.), anhydrous potassium carbonate (1.26 g, 9.1 mmol, 2 equiv.) and MeOH (20 mL) were added and the mixture was refluxed overnight under an argon atmosphere. After cooling to RT, TBME (12 mL) and water (12 mL) were added, the organic phase was separated and the aqueous phase extracted with additional TBME (2 x 25 mL). The combined organic layers were washed with brine, dried (Na₂SO₄) and concentrated to give a residue that was purified by flash chromatography on silica gel using 5-10% EtOAc in hexane as eluent: 1.21 g (79%).

1-Cyclopentyl-3-methyl-2-(2-pyridyl)indole-5-carboxylic acid:

The methyl ester from above (1.10 g, 3.29 mmol) was dissolved in 1 : 1 THF-MeOH (10 mL) and 6 N aqueous NaOH (4.2 mL, 5 equiv.) was added. The mixture was stirred for 3 h at 60 °C. The reaction mixture was cooled to RT and water (15 mL) was added. The suspension was filtered and the yellow filtrate washed with TBME (2 x 20 mL) and hexane (20 mL). Conc. HCl (1.9 mL) was added dropwise with stirring to pH 8, followed by AcOH (3 mL) to pH 2. The suspension was cooled and the white solid collected by filtration. The product was then dried under vacuum: 0.90 g (86%).

EXAMPLE 4

3-Cyclohexyl-2-furan-3-yl-7-methyl-1H-indole-6-carboxylic acid methyl ester



Step 1: 3-Amino-2-methylbenzoic acid (15.00 g, 0.099 mol) was suspended in MeOH (150 mL) and thionyl chloride (25.33 mL, 0.347 mole, 3.5 equiv.) was added dropwise. The mixture was heated overnight at 70 °C. After cooling to RT, volatiles were removed under reduced pressure and the residue triturated with ether (150 mL). The solid was filtered off and dried (18.36 g). The solid was suspended in DCM (600 mL) and saturated aqueous NaHCO₃ (250 mL) was added. After stirring for 15 minutes, the organic layer was separated and washed successively with NaHCO₃ solution (2 x 250 mL), water (250 mL) and brine (250 mL). The solution was dried (Na₂SO₄), filtered and evaporated to dryness to give the desired aniline (14.8 g, 90% yield).

Steps 2 and 3: The ester obtained in the previous step (12.50 g, 75.6 mmol) was dissolved in DCM (190 mL) and methylthioacetaldehyde dimethyl acetal (10.1 mL, 75.6 mmol) was added. The solution was cooled to -30 °C. N-chlorosuccinimide (10.10 g, 75.6 mmol) was added in 6 portions over 30 minutes. Triethylamine (10.6 mL, 75.6 mmol) was then added dropwise over 10 min and after stirring for an additional 15 min, the cooling bath was removed and the temperature brought to reflux. After 5 h, the reaction mixture was cooled to RT and evaporated to dryness. The residue was dissolved in ether (750 mL) and 2M HCl (303 mL) was added. After stirring at RT for 1.5 h, the ether layer was separated and washed with NaHCO₃ solution (2 x 150 mL) and brine (250 mL). The original acidic aqueous

phase was extracted with DCM (2 x 100 mL) and the extracts washed as above and then combined with the first ether phase. The combined organic phases were dried (Na_2SO_4) and evaporated to dryness and the material purified by flash chromatography on silica gel using 30-0% hexane in DCM as eluent to give the
5 desired 3-thiomethylindole derivative (9.37 g).

Step 4: The thiomethyl indole obtained in the previous step (8.37 g, 35.4 mmol) was dissolved in absolute EtOH (220 mL) and Raney-nickel (Ra-Ni, 25 g) was added. After stirring at RT for 3 h, another portion of Ra-Ni (15 g) was added and stirring
10 resumed for an additional 45 min. The mixture was filtered and the filtrate evaporated to dryness to give the desired indole (6.26 g, 93%).

Step 5: The indole ester obtained in the previous step (4.00 g, 21 mmol) was dissolved in a mixture of MeOH (18 mL) and water (18 mL). KOH (11.86 g, 210
15 mmol) was added and the mixture stirred at 75 °C for 2 h. Cyclohexanone (7.26 g, 74 mmol, 3 equiv.) was added dropwise over 15 min and stirring at 75 °C was continued overnight. MeOH was removed under reduced pressure and water (500 mL) was added to the residue. Insoluble material was removed by filtration and the aqueous phase was then washed with TBME (200 mL). The aqueous phase was
20 acidified to pH 4 with formic acid to produce a white precipitate that was collected by filtration, washed with water and dried. The desired cyclohexenylindole was obtained (4.77 g, 89%).

Step 6: The cyclohexene derivative from above (4.70 g, 18 mmol) was dissolved in
25 MeOH (1 L) and 20% $\text{Pd}(\text{OH})_2$ on C (0.24 g) was added. The mixture was hydrogenated (60 psi of H_2 gas) for 7 h, filtered, and then concentrated under reduced pressure. The residue was triturated with MeOH (100 mL), and the solid collected by filtration and dried (3.50 g)

Step 7: The carboxylic acid from step 6 (3.44 g, 13 mmol) was dissolved in DMF (22 mL) and anhydrous K_2CO_3 (2.55 g, 18 mmol) was added followed by iodomethane (2.09 g, 15 mmol). The mixture was stirred overnight at room temperature and then poured into water (60 mL) and acidified to pH 4 with 4 N HCl (10 mL). The product was extracted with ether (3 x 150 mL) and washed with water (4 X 90 mL) and brine
35 (90 mL). After drying (Na_2SO_4) and removal of solvents, the desired methyl ester


was obtained as a white solid (3.44 g).

Step 8: The indole derivative of step 7 (3.20 g, 12 mmol) was dissolved in a mixture of THF (30 mL) and CHCl_3 (30 mL) and the solution cooled to $-5\text{ }^\circ\text{C}$. Pyridinium
5 tribromide (5.17 g, 16 mmol) was added in portions maintaining the internal temperature at $-5\text{ }^\circ\text{C}$. After stirring for 1 h at $-5\text{ }^\circ\text{C}$, the reaction was allowed to warm up to room temperature and concentrated to dryness. Water (50 mL) was added and the product extracted with ether (3 x 100 mL). The extract was washed with water (2 X 90 mL) and brine (90 mL) and dried (Na_2SO_4). Removal of solvents
10 under reduced pressure gave a residue that was purified by flash chromatography using 10% EtOAc in hexane as eluent. The desired bromide was obtained as a solid (3.23 g).

Step 9: The bromoindole of step 8 (3.10 g, 8.9 mmol) was dissolved in a mixture of
15 toluene (20 mL), EtOH (20 mL) and water (14 mL). 3-Furylboronic acid (1.27 g, 11.3 mmol), LiCl (0.75 g, 17.7 mmol) and Na_2CO_3 (2.35 g, 22.1 mmol) were added. The solution was degassed by bubbling argon gas for 30 min and tetrakis(triphenylphosphine)palladium⁰ (0.41 g, 0.35 mmol) was added. The mixture was heated to $80\text{ }^\circ\text{C}$ for 3 h, cooled to room temperature and extracted with TBME
20 (350 mL) using water (90 mL) for rinsing undissolved solids. The organic phase was separated and the aqueous phase extracted twice more with TBME (2 X 150 mL). The combined extracts were washed with water (2 X 100 mL) and brine (100 mL), dried (Na_2SO_4) and concentrated to dryness. The residue was purified by flash chromatography using 5-15% EtOAc in hexane as eluent (3.0 g).
25 The protected indole from above can be saponified to the free carboxylic acid as described in example 3.

EXAMPLE 5:

30 **Methyl 2-bromo-3-cyclohexyl-1H-indole-5-carboxylate and 3-cyclohexyl-2-furan-3-yl-1H-indole-5-carboxylic acid:**




Using the same series of reactions described in example 2 of WO 03/010141 but starting from indole 5-carboxylic acid, methyl 2-bromo-3-cyclohexyl-1H-indole-5-carboxylate was obtained and elaborated to 3-cyclohexyl-2-furan-3-yl-1H-indole-5-carboxylic acid using protocols such as those described in examples 3 and 4.

EXAMPLE 6:

7-Methoxy-1H-indole-6-carboxylic acid methyl ester:

10



Step 1: 4-Methylsalicylic acid (32.1 g, 0.21 mol) and potassium carbonate (61.2 g, 0.44 mol) were suspended in acetone (300 mL) and the mixture brought to reflux temperature. Dimethyl sulfate (66.5 g, 0.53 mol) was added dropwise within 1 h and stirring continued overnight at reflux. Additional dimethylsulfate (30 mL) and potassium carbonate (2 x 15 g) were added and refluxing for an additional 24 h was required to complete the reaction. The reaction mixture was then cooled to room temperature and inorganic salts removed by filtration using acetone for washings. The filtrate was evaporated under reduced pressure and the oily residue was dissolved in MeOH (300 mL). Concentrated ammonium hydroxide (90 mL) was added and the mixture was stirred for 30 minutes at room temperature. Methanol was removed in vacuo and the residue portioned between ether (300 mL) and water (200 mL). The organic phase was separated and washed with brine and dried (Na₂SO₄). Evaporation of the ether gave the desired di-methylated product as a yellow oil (38.1 g) that was used directly in the next step.

Step 2: The ester obtained in the previous step (38.0 g, 0.21 mol) was dissolved in AcOH (250 mL) and bromine (37.2 g, 0.23 mole, 1.1 equiv.) was added dropwise over 30 min with stirring at room temperature. After completion, the reaction mixture
5 was stirred for an additional hour, at which point TLC analysis indicated complete conversion. The reaction mixture was poured into water (1 L) and solid Na₂CO₃ was added cautiously with stirring until the mixture was neutral. The off-white precipitate that formed was collected by filtration, washed with water and dried to give the desired bromo derivative (47.2 g).

10

Step 3: The bromo derivative obtained in the previous step (44.5 g, 0.17 mol) was added in small portions to conc. H₂SO₄ (170 mL) and the mixture was stirred in an ice-salt bath until all solids dissolved. Conc. HNO₃ (17 mL) was then added dropwise over 20 min and stirring continued for an additional hour in the ice bath.
15 The reaction mixture was then slowly added to ice-water (2 L) and the precipitated yellow solid was collected by filtration. The solid was washed with water, NaHCO₃ solution and water again. After drying, the desired nitro derivative was obtained as an orange solid (36.8 g).

20 **Step 4:** The product obtained in the previous step (129.0 g, 0.42 mol) was dissolved in DMF (400 mL) and DMF-dimethyl acetal (151.6 g, 1.27 mole, 3 equiv.) was added in one portion. The mixture was heated at 110-120 °C under an argon atmosphere until conversion was judged complete by TLC (24 h). The reaction mixture was cooled to room temperature and volatiles removed under vacuum to give a dark
25 colored residue (~180 g). Trituration from ether-THF gave the desired enamine as red crystals (72 g).

Step 5: The enamine obtained in the previous step (72.0 g, 0.20 mol) was dissolved in a mixture of THF (600 mL) and MeOH (600 mL). The dark red solution was
30 heated to 30 °C and Raney-Nickel (18 g) was added to the solution. Hydrazine hydrate (11.6 g, 0.23 mole, 1.15 equiv.) was then added dropwise over 30 min. The reaction temperature was increased to 50 °C and a second portion of hydrazine hydrate (11.6 g, 0.23 mole, 1.15 equiv.) was added over 30 min. After stirring overnight at 50 °C, additional Raney-nickel (20 g) and hydrazine hydrate (11.6 g,
35 0.23 mole, 1.15 equiv.) were added and after stirring for another 7 h at 50 °C, the

reaction was judged complete by TLC. After cooling, the catalyst was removed by filtration through a pad of celite and the filtrate was evaporated under reduced pressure. The dark brown residue was dissolved in EtOAc (3 L) and the solution washed with water (1.5 L), 10% HCl (1 L) and brine (700 mL). After drying (Na_2SO_4),
5 removal of solvents gave the desired bromoindole derivative as a brown solid (35 g).

Step 6: The bromoindole derivative obtained in the previous step (35 g) was dissolved in MeOH (1 L) and triethylamine (16.3 g, 1.2 equiv.) was added followed by 10% Pd/C (1.06 g). The mixture was stirred under hydrogen (35 psi) until completion
10 of the reaction (7 h). The catalyst was then removed by filtration and volatiles removed under reduced pressure. The residue was dissolved in EtOAc (700 mL) and the solution washed with 10% HCl (300mL), water (350 mL), NaHCO_3 (350 mL) and brine. The solution was dried (Na_2SO_4) and concentrated under reduced pressure to give the desired indole as a light brown solid (25 g).

15

This indole derivative was saponified under standard conditions and elaborated to final inhibitors as described in examples 3 and 4.

EXAMPLE 7

20 Inhibition of NS5B RNA dependent RNA polymerase activity

The compounds of the invention were tested for inhibitory activity against the hepatitis C virus RNA dependant polymerase (NS5B), according to the protocol described in WO 03/010141.

25 EXAMPLE 8

Specificity of NS5B RNA dependent RNA polymerase inhibition

The compounds of the invention were tested for inhibitory activity against polio virus RNA dependent RNA polymerase and calf thymus DNA dependent RNA polymerase II in the format that is described in WO 03/010141 for the HCV polymerase, with the
30 exception that another polymerase was used in place of the HCV NS5B polymerase.


In Tables 1 and 2 below, the following ranges apply:

IC_{50} : A = 20 μM -1 μM and B < 1 μM

35




























The following compounds are prepared, defined by A, B, R², R³ and Z according to the table 1.




























TABLE 1





















5

Cpd. #	A	B	R ²	R ³	Z	IC ₅₀	MS (MH ⁺)
101	-N(CH ₃)-	=C-				B	475.1
102	=N-	-N-				A	447.1
103	=N-	-N-				A	461.1
104	=N-	-N-				A	399.1
105	=N-	-N-				A	521.2


Cpd. #	A	B	R ²	R ³	Z	IC ₅₀	MS (MH ⁺)
106	=N-	-N-				A	385.1
107	=N-	-N-				B	507.1
108	=N-	-N-				B	136.1
109	=N-	-N-				A	374.1
110	=N-	-N-				B	496.1
111	=N-	-N-				B	450.1
112	=N-	-N-				B	388.1
113	=N-	-N-				B	510.1
114	-N(CH ₃)-	=C-				B	463.1

Cpd. #	A	B	R ²	R ³	Z	IC ₅₀	MS (MH ⁺)
115	-N(CH ₃)-	=C-				B	413.1
116	-N(CH ₃)-	=C-				B	441.2
117	-N(CH ₃)-	=C-				B	461.1
118	=C(CH ₃)-	-N-				B	449.1
119	=C(CH ₃)-	-N-				A	398.1
120	-O-	=C-				A	414.1
121	-S-	=C-				B	390.1
122	-S-	=C-				A	427.1
123	-N(CH ₃)-	=C-				B	398.1

Cpd. #	A	B	R ²	R ³	Z	IC ₅₀	MS (MH ⁺)
124	-NH-	=C-				B	449.2
125	-NH-	=C-				B	373.1
126	-N(CH ₃)-	=C-				A	558.3
127	=C(CH ₃)-	-N-					
128	-O-	=C-					
129	-N(CH ₃)-	=C-					

The following compounds are prepared, defined by R², R³, R^{4a} and Z according to the table 2. The position number p being 2 or 3 indicates to which C-atom the group C(=O)-Z is bonded.

TABLE 2



Cpd. #	R ²	R ³	R ^{4a}	p	Z	IC ₅₀	MS (MH ⁺)
201			-OCH ₃	2		B	417.1
202			-OCH ₃	2		B	479.1
203			-H	3		A	401.1
204			-H	3		A	463.1